

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Magellan[®] 800WG

Systemic Fungicide

ACTIVE CONSTITUENT: **800 g/kg FOSETYL-ALUMINIUM**

GROUP

33

FUNGICIDE

For the prevention and control of Phytophthora rots in ornamentals and Pythium in turf as specified in the Direction for Use

Formulation type
Water Dispersible
Granule

WG

 **CENTAUR**
ASIA PACIFIC
www.centaur-asiapacific.com
info@centaur-asiapacific.com



adama.com

NET CONTENTS: 1 - 25 kg

DIRECTIONS FOR USE

RESTRAINT

DO NOT use as an alternative to steam sterilisation of potting soils used in nurseries.

SITUATION	DISEASE	RATE	CRITICAL COMMENTS
Ornamentals	Crown and Root Rot (<i>Phytophthora</i> spp.)	90 g/100 L water (10 g/m ²)	Apply as a soil drench at intervals of 6 weeks. Container grown plants: Drench volume depends on size. For 150 mm container apply 200 mL.
Turf (golf and bowling greens and other intensely managed Turf)	<i>Pythium</i> spp.	125 g/100 m ²	Begin preventative applications when conditions first favour disease and continue as long as conditions are favourable for disease at approximately 21 day intervals. Apply as a foliar spray. DO NOT mow or water treated area until foliage is completely dry.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

GENERAL INSTRUCTIONS

MIXING

Fill the spray tank to at least half the desired amount with clean water and commence agitation. Add the required quantity of MAGELLAN® 800WG Systemic Fungicide directly to the water and complete filling to the final volume. Continue agitation. DO NOT make a slurry or paste prior to adding to the tank.

APPLICATION

Turf only: Apply 5 - 20 L of water per 100 m².

CROP SAFETY

Ornamentals

MAGELLAN® 800WG Systemic Fungicide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical products. It is advisable to treat only a small number of plants first, in order to determine their reaction to the product, before large scale use.

COMPATIBILITY

MAGELLAN® 800WG Systemic Fungicide is compatible with a number of fungicides. If unsure, it is recommended that a Jar test be conducted to determine compatibility. Always add MAGELLAN® 800WG Systemic Fungicide to the mixture last. With any mixture, constantly agitate prior to and during application. It is not recommended to mix this product with more than one additional pesticide in the tank. Always test mixtures on a small number of plants before large scale use. As formulations of other manufacturers products are beyond the control of Adama Australia, all mixtures should be tested prior to mixing commercial quantities.

RESISTANCE WARNING

MAGELLAN® 800WG

Systemic Fungicide is a member of the Phosphonates group of fungicides. For fungicide resistance management MAGELLAN® 800WG Systemic Fungicide is a Group 33 Fungicide. Some naturally occurring fungi resistant to MAGELLAN® 800WG Systemic Fungicide or any other Group 33 Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. The resistant fungi will not be controlled by MAGELLAN® 800WG Systemic Fungicide or other Group 33 Fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Adama Australia accepts no liability for any losses that may result from the failure of MAGELLAN® 800WG Systemic Fungicide to control resistant fungi.



PROTECTION OF LIVESTOCK

DO NOT graze treated turf or lawn; or feed turf or lawn clippings from any treated area to poultry or livestock.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN. Store in closed original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. Avoid inhaling dust or spray mist. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for MAGELLAN® 800WG Systemic Fungicide is available from adama.com or call Customer Service on 1800 423 262.

CONDITIONS OF SALE: The use of MAGELLAN® 800WG Systemic Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Adama Australia regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Adama Australia accepts no responsibility for any consequence whatsoever from the use of this product.

® Registered trademark of an Adama Group Company

10/2017 9441