

TECHNICAL NOTES

Date: 02/04/25 Issue Number: 6

ACTIVE INGREDIENT: FLUOPYRAM

EFFECTIVE STING NEMATODE MANAGEMENT WITH INDEMNIFY

The Southern Sting Nematode (*Ibipora Iolii*) is one of the most aggressive and damaging nematodes in turfgrass, particularly in sandy soils. It affects high-value turf like golf courses, sports fields, and turf farms, leading to severe turf decline.

→ Biology & Life Cycle

Large, slender nematode with a spear-like stylet for feeding.

Prefers sandy soils (>80% sand), which are common in many turf situations.

→ Life Cycle

Egg \rightarrow Four Juvenile Stages (J1–J4) \rightarrow Adult.

Requires both males and females for reproduction.

Most active during warmer months (spring to autumn) when soil temperatures are optimal.

→ Feeding & Damage

Ectoparasitic feeder, meaning it stays outside the root and inserts its stylet to extract nutrients.

Causes "sting" symptoms – stunted roots, root tip swelling, and necrosis.

Turf becomes thinned, chlorotic, and weakens due to poor root function.

Damage is worse in sandy soils with low organic matter, common in coastal golf courses, sports fields, and amenity turf in Australia.

Population levels peak in warm seasons and decline in winter.

ABOUT INDEMNIFY

Indemnify® is a proven, easy-to-use nematicide that controls key plant parasitic nematodes and improves root growth and turfgrass health resulting in better playability with improved wear tolerance.

→ Targeted Application Benefits

Indemnify contains the active ingredient Fluopyram. Fluopyram is a succinate dehydrogenase inhibitor (SDHI) that paralyses and eventually kills nematodes by blocking cellular respiration.

Fluopyram is a contact nematicide with acropetal (upwards) movement.


The effect on nematodes is initial paralysis and this effect begins once the nematicide makes contact. In general, visual improvement to damaged turf occurs four weeks after application.

Effects of improved turfgrass quality can last between two to six months. The exact duration is dependent on several factors including, but not limited to, nematode species and turfgrass type.

First standard parties projection on comments of the comments

Results run deeper with Indemnify

- + Reduced summer hand watering
- + More summer aeration
- + Reduced root diseases
- + Improved surface quality
- + Better surface stability







TECHNICAL NOTES

APPLICATION STRATEGY

Indemnify should be applied as a preventative treatment when conditions are favourable for root growth, before nematode populations build up to damaging levels. The population buildup of some key species such as sting nematode often occurs during or following a period of root growth, so optimum timing of application, while situation-dependent, will often be just prior to periods of peak root growth. This timing is typically early spring and autumn.

Apply up to four applications at the higher curative spot treatment rate if controlling outbreaks that are severe or are expected to become severe.

TREATMENT TIPS FOR BEST RESULTS

- Apply in 400 800 L water/ha.
- Apply 0.625L/ha as a preventative treatment.
- Apply 1.25L/ha as a curative spot treatment.
- Apply no more than 4 applications a year refer to <u>label</u> for application rates.

→ Tank mixing


Do not tank mix this product with soil wetting agents or surfactants.

The <u>Indemnify</u> formulation contains soil penetration adjuvants to optimise delivery of the active ingredient to the root zone.



INDEMNIFY TRIAL RESULTS

<u>Indemnify</u> trial showing reduction in sting nematodes for various treatment programmes.



STEWARDSHIP

Always read the label before use.

For full list of application and usage recommendations, view the <u>label</u>.

CONTACT US

For further information on controlling Sting Nematodes with Indemnify, please contact the Envu team.

. Details can be found on the Envu website by visiting <u>Our Team</u> in Australia.

Find out more about Indemnify



Distributed by:



www.centaur-asiapacific.com info@centaur-asiapacific.com

*Singapore only.

